



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned} s+a &= y^2 + 2yz + z^2 + x(2x+y+z) = (y+z)^2 + x^2 = 2x^2; \\ \text{similarly, } s+b &= 2y^2; \\ s+c &= 2z^2. \end{aligned}$$

Hence $\sqrt{s+a} + \sqrt{s+b} + \sqrt{s+c} = (x+y+z)\sqrt{2} = 0$, an equation involving only a, b, c .

If we wish to express this relation without radicals we transpose, square, and reduce; whence $a-s = \sqrt{(s+b)(s+c)}$, whence

$$\begin{aligned} s^2 + as &= a^2 - bc \\ s^2 + bs &= b^2 - ca \\ s^2 + cs &= c^2 - ab. \end{aligned}$$

Whence $4s^2 = a^2 + b^2 + c^2 - (ab + bc + ca)$; and therefore

$$a^2 + b^2 + c^2 + 3(ab + bc + ca) = 0.$$

This may also be written

$$\frac{a}{bc} + \frac{b}{ca} + \frac{c}{ab} + 3\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right) = 0.$$

Example. Let $x = +1, y = +2, z = -3$; then $a = -5, b = +1, c = +11$.

Excellent solutions of this problem were received from *F. L. SAWYER, LON C. WALKER, JOSIAH H. DRUMMOND, J. K. ELLWOOD, J. SCHEFFER*, and *G. B. M. ZERR*.

Mr. Baker sent in neat solutions of problems 151 and 152.

154. Proposed by *F. P. MATZ, Sc. D., Ph. D.*, Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

Show that the equation, $x^4 + qx^2 + s = 0 \dots (1)$, can not have three *equal* roots.

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy in Ohio University, Athens, Ohio.

By the usual theory, the conditions that $ax^4 + 4bx^3 + 6cx^2 + 4dx + e = 0 \dots (1)$ shall have three equal roots are

$$ae - 4bd + 3c^2 = 0 \dots (2),$$

$$ad^2 + eb^2 + c^3 - ace - 2bcd = 0 \dots (3).$$

In the given equation, $x^4 + qx^2 + s = 0 \dots (4)$, $a = 1, b = 0, c = \frac{1}{2}q, d = 0, e = s$, and (2) and (3) become

$$s + \frac{q^2}{12} = 0 \dots (5), \quad \frac{q^3}{36} - qs = 0 \dots (6).$$

Eliminating s from (5) and (6) gives $q=0$, which is inconsistent with the supposition.

Also solved similarly by *G. B. M. ZERR, F. L. SAWYER, JOSIAH H. DRUMMOND, J. R. HITT, and HARRY S. VANDIVER.*

155. Proposed by *WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.*

If the roots of the cubic $x^2 + 3px^2 + 3qx + r = 0$ be in harmonical progression, $2q^3 = r(3pq - r)$.

Solution by *HOMER R. HIGLEY, M. S., State Normal School, East Stroudsburg, Pa.; J. R. HITT, Coronal Institute, San Marcos, Tex.; H. S. VANDIVER, Bala, Pa.; and the PROPOSER.*

If α, β, γ be the roots of a cubic, we have

$$x^3 - (\alpha + \beta + \gamma)x^2 + (\alpha\beta + \alpha\gamma + \beta\gamma)x - \alpha\beta\gamma = 0 \dots (1).$$

By the condition of the problem,

$$\beta = \frac{2\alpha\gamma}{\alpha + \gamma}, \text{ or } \alpha\beta + \alpha\gamma + \beta\gamma = 3\alpha\gamma \dots (2),$$

and (1) becomes

$$x^3 - (\alpha + \beta + \gamma)x^2 + 3\alpha\gamma x - \alpha\beta\gamma = 0 \dots (3).$$

Comparing (3) and

$$x^3 + 3px^2 + 3qx + r = 0 \dots (4),$$

$$\alpha + \beta + \gamma = -3p \dots (5), \quad \alpha\gamma = q \dots (6), \quad \text{and } \alpha\beta\gamma = -r \dots (7).$$

From (6) and (7), $\beta = -r/q \dots (8)$.

From (5) and (8), $\alpha + \gamma = r/q - 3p \dots (9)$.

Eliminating α, β, γ from (2) by use of (6), (8), (9), $2q^3 = r(3pq - r)$.

Also solved by *J. SCHEFFER, LON C. WALKER, JOSIAH H. DRUMMOND, F. L. SAWYER, and G. B. M. ZERR.*

GEOMETRY

183. Proposed by *S. F. NORRIS, Professor of Mathematics and Astronomy in Baltimore City College, Baltimore, Md.*

Two quadrilaterals having three sides of the one equal to the three corresponding sides of the other, each to each, and the two corresponding angles adjacent to the unknown sides equal, each to each, are equal figures. [*Olney's Geometry*, page 129.]

This is problem 169, an erroneous demonstration of which was published in the April number of the current volume of the MONTHLY. The fallacy in the demonstration is pointed out in Dr. Halsted's article, "Proving the False," on page 129. Ed.